

Peer reviewed scientific journals (*Equal contribution, +supervised student)

Gavish Y, Marsh CJ, Kuemmerlen M, **Stoll S**, Haase P, Kunin WE (2017) Accounting for biotic interaction through alpha-diversity constraints in stacked species distribution models. *Methods in Ecology and Evolution* 8: 1092–1102.

Bowler DE, Hof C, Haase P, Kröncke I, Schweiger O, Adrian R, Baert L, Bauer H-G, Blick T, Brooker RW, Dekoninck W, Domisch S, Eckmann R, Hendrickx F, Hickler T, Klotz S, Kraberg A, Kühn I, Matesanz S, Meschede A, Neumann H, O'Hara B, Russell, DJ, Sell AF, Sonnewald M, **Stoll S**, Sundermann A, Tackenberg O, Türkay M, Valladares F, van Herk K, van Klink R, Vermeulen R, Voigtländer K, Wagner R, Welk E, Wiemers M, Wiltshire KH, Böhning-Gaese K (2017) Cross-realm assessment of climate-change impacts on species' abundance trends. *Nature Ecology and Evolution*: 67.
doi:10.1038/s41559-016-0067.

Stoll S, +Hormel N, +Früh D, Tonkin JD (2017) Effects of *Chaetogaster limnaei limnaei* on freshwater snail communities. *Hydrobiologia* 785: 101-113.

+Früh D, Haase P, **Stoll S** (2017) Temperature drives asymmetric competition between alien and indigenous freshwater snail species, *Physa acuta* and *Physa fontinalis*. *Aquatic Sciences* 79: 187–195.

Li F, Sundermann A, **Stoll S**, Haase P (2016) A newly developed dispersal capacity metric indicates succession of benthic invertebrates in restored rivers. *Science of the Total Environment* 569: 1570-1578.

Kuemmerlen M, **Stoll S**, Sundermann A, Haase P (2016) Predicting stream macroinvertebrate distributions in the LTER-site Rhine-Main-Observatory: Long-term monitoring data meets high-resolution, catchment-based SDMs. *Ecological Indicators* 65: 122-132.

Haase P, Frenzel M, Klotz S, Musche M, **Stoll S** (2016) The Long-Term Ecological Research (LTER) network: Relevance, current status, future perspective and examples from marine, freshwater and terrestrial long-term observation. *Ecological Indicators* 65: 1-3.

Stoll S, +Breyer P, Tonkin JD, +Früh D, Haase P (2016) Scale-dependent effects of river habitat quality on benthic invertebrate communities - implications for stream restoration. *Science of the Total Environment* 553: 495-503.

Tonkin JD, **Stoll S**, Jähnig S, Haase P (2016) Elements of metacommunity structure of river and riparian assemblages: communities, taxonomic groups and deconstructed trait groups. *Ecological Complexity* 25: 35-43.

Müller F, Bergmann M, Dannowski R, Dippner JW, Gnauck A, Haase P, Jochimsen MC, Kasprzak P, Kröncke I, Kümmerlin R, Küster M, Lischeid G, Meesenburg H, Merz C, Millat G, Müller J, Padisák J, Schimming CG, Schubert H, Schult M, Selmečzy G, Shatwell T, **Stoll S**, Schwabe M, Soltwedel T, Straile D (2016) Assessing resilience in long-term ecological data sets. *Ecological Indicators* 65: 10-43.

Tonkin JD, **Stoll S**, Jähnig S, Haase P (2016) Anthropogenic stress alters community concordance at the river-riparian interface. *Ecological Indicators* 65: 133-141.

Tonkin JD, **Stoll S**, Jähnig S, Haase P (2016) Contrasting metacommunity structure and beta diversity in an aquatic-floodplain system. *Oikos* 125: 686–697.

Geijzendorffer IR, Regan E, Pereira HM, Brotons L, Brummitt N, Gavish Y, Haase P, Martin CS, Mihoub J-B, Secades C, Schmeller DS, **Stoll S**, Wetzel FT, Walters M (2016) Bridging the gap between biodiversity data and policy reporting needs: An Essential Biodiversity Variables perspective. *Journal of Applied Ecology* 53: 1341–1350.

Simaika JP*, **Stoll S***, Lorenz AW, Thomas G, Sundermann A, Haase P (2015) Bundles of stream restoration measures and their effects on fish communities. *Limnologica* 55: 1-8.

+Höckendorff S, +Früh D, +Hormel N, Haase P, **Stoll S** (2015) Biotic interactions under climate warming: temperature-dependent and species-specific effects of the oligochaete *Chaetogaster limnaei* on snails. *Freshwater Science* 34(4): 1304-1311.

Thomas G, Lorenz AW, Sundermann A, Haase P, Peter A, **Stoll S** (2015) Fish community responses and the temporal dynamic of recovery following river habitat restorations in Europe. *Freshwater Science* 34(3): 975-990.

Domisch S, Jähnig SJ, Simaika JP, Kuemmerlen M, **Stoll S** (2015) Application of species distribution models in stream ecosystems: the challenge of spatial scale, environmental predictors and species occurrence data. *Fundamental and Applied Limnology* 186(1-2): 45-61.

Stoll S, Frenzel M, Burkhard B, Adamescu M, Augustaitis A, Baeßler C, Bonet FJ, Cazacu C, Cosor GL, Díaz-Delgado R, Carranza ML, Grandin U, Haase P, Hämäläinen H, Loke R, Müller J, Stanisci A, Staszewski T, Müller F (2015) Assessment of ecosystem integrity and service gradients across Europe using the LTER Europe network. *Ecological Modelling* 295: 75-87.

Tonkin JD, **Stoll S**, Sundermann A, Haase P (2014) Dispersal distance and taxonomic pool, not barriers, delimit colonisation of restored river reaches by benthic invertebrates. *Freshwater Biology* 59: 1843–1855.

Stoll S, Kail J, Lorenz AW, Sundermann A, Haase P (2014) The importance of the regional species pool, ecological species traits and local habitat conditions for the colonization of restored river reaches by fish. *PLoS ONE* 9(1): e84741. doi:10.1371/journal.pone.0084741

Lorenz AW, **Stoll S**, Sundermann A, Haase P (2013) Do adult and YOY fish benefit from river restoration measures? *Ecological Engineering* 61: 174-181.

Stoll S, +Früh D, +Westerwald B, +Hormel N, Haase P (2013) Density-dependent relationship between *Chaetogaster limnaei* and the freshwater snail *Physa acuta* (Pulmonata). *Freshwater Science* 32: 642-649.

Stoll S (2013) Interactive effect of trophic state and spring water level increase on commercial bream (*Abramis brama*) yields in Lake Constance. *Journal of Applied Ichthyology* 29: 617-622.

Stoll S, Sundermann A, Lorenz AW, Kail J, Haase P (2013) Small and impoverished fish species pools are a main challenge to the colonization of restored river reaches. *Freshwater Biology* 58: 664-674.

+Früh D, **Stoll S**, Haase P (2012) Physico-chemical variables determining the invasion risk of freshwater habitats by alien mollusks and crustaceans. *Ecology and Evolution* 2(11): 2843–2853. doi: 10.1002/ece3.382.

Stoll S, Beeck P (2012) Larval fish in troubled waters - is the behavioural response of larval fish to hydrodynamic stress active or passive? *Canadian Journal of Fisheries and Aquatic Sciences* 69(10): 1576-1584.

+Früh D, **Stoll S**, Haase P (2012) Physicochemical and morphological degradation of stream and river habitats increases invasion risk. *Biological Invasions* 14: 2243-2253.

Kappes H, **Stoll S**, Haase P (2012) Behavioral differences between native species and an invader (Gastropoda: *Arion vulgaris*). *Belgian Journal of Zoology* 142: 49-58.

Stoll S, Beeck P (2011) Post-release stranding rates of stocked larval allis shad (*Alosa alosa*) larvae exposed to surface waves. *Journal of Applied Ichthyology* 27 (Suppl. 3): 41-44.

Sundermann A*, **Stoll S***, Haase P (2011) River restoration success depends on the species pool in the surroundings *Ecological Applications* 21: 1962-1971.

Stoll S, Fischer P (2011) Three different patterns how low-intensity waves can affect growth of littoral fish: a mesocosm study. *Oecologia* 165: 567-576.

Gabel F*, **Stoll S***, Fischer P, Pusch M, Garcia X-F (2011) Waves impact predator-prey interactions between fish and invertebrates: an experimental study. *Oecologia* 165: 101-109. *The first two authors contributed equally to this work.

Stoll S, Hofmann H, Fischer P (2010) Effect of wave exposure dynamics on gut content mass and growth of young-of-the-year fishes in the littoral zone of lakes. *Journal of Fish Biology* 76: 1714-1728.

Stoll S, Probst WN, Eckmann R, Fischer P (2010) A mesocosm experiment investigating the effects of substratum quality and wave exposure on the survival of fish eggs. *Aquatic Sciences* 72: 509-517.

Probst WN, **Stoll S**, Peters L, Fischer P, Eckmann R (2009) Lake water level increase during spring affects the breeding success of bream (*Abramis brama*). *Hydrobiologia* 632: 211-224.

Probst WN, **Stoll S**, Hofmann H, Fischer P, Eckmann R (2009) Spawning site selection by Eurasian perch (*Perca fluviatilis* L.) in relation to temperature and wave exposure. *Ecology of Freshwater Fish* 18: 1-7.

Stoll S, Fischer P, Klahold P, Scheifhacken N, Hofmann H, Rothhaupt K-O (2008) Effects of water depth and hydrodynamics on the growth and distribution of juvenile cyprinids in the littoral zone of a large prealpine lake. *Journal of Fish Biology* 72: 1001-1022.

Non-peer reviewed publications:

Tonkin JD, **Stoll S**, Jähnig S, Haase P (2015) Variable elements of metacommunity structure across an aquatic-terrestrial ecotone. PeerJ PrePrints 3:e1261. [dx.doi.org/10.7287/peerj.preprints.1026v1](https://doi.org/10.7287/peerj.preprints.1026v1).

Früh D, Haase P, **Stoll S** (2015) Temperature drives asymmetric competition between alien and indigenous freshwater snails (*Physa acuta* vs. *Physa fontinalis*). PeerJ PrePrints 3:e1227 [dx.doi.org/10.7287/peerj.preprints.997v1](https://doi.org/10.7287/peerj.preprints.997v1).

Li F, Sundermann A, **Stoll S**, Haase P (2015) A newly developed dispersal capacity metric indicates succession of benthic invertebrates in restored rivers. PeerJ PrePrints 3:e1125 [dx.doi.org/10.7287/peerj.preprints.911v1](https://doi.org/10.7287/peerj.preprints.911v1).

Tonkin JD, **Stoll S**, Jähnig SC, Haase P (2015) Anthropogenic stress alters community concordance at the river-riparian interface. PeerJ PrePrints 3:e798v1 [dx.doi.org/10.7287/peerj.preprints.798v1](https://doi.org/10.7287/peerj.preprints.798v1).

Ostendorp W, Brünner H, Dienst M, Eckmann R, Fiebrich M, Fiedler W, Gabel F, Grabher M, Heck K, Hermann M, Jacoby H, Kiechle J, Knötzsch G, Kramer I, Lehmann A, Löderbusch W, Miler O, Müller M, Rieder J, Sättele B, Schäfer C, Schmidt B, Schmitz G, Schürenberg B, **Stoll S**, Trautner J, Wilhelm P, Winkler U (2014) Effects of lake bank retaining walls at Lake Constance/Untersee (Germany, Switzerland) on the littoral biota: results of a scenario-based expert judgement. *Mitteilungen des Badischen Landesvereins für Naturkunde und Naturschutz* 21(3): 371 - 404.

Ostendorp W, Brünner H, Dienst M, Eckmann R, Fiebrich M, Fiedler W, Gabel F, Grabher M, Heck K, Hermann M, Jacoby H, Kiechle J, Knötzsch G, Kramer I, Lehmann A, Löderbusch W, Miler O, Müller M, Rieder J, Sättele B, Schäfer C, Schmidt B, Schmitz G, Schürenberg B, **Stoll S**, Trautner J, Wilhelm P, Winkler U (2015) Ökologische Auswirkungen von Ufervorschüttungen am Bodensee-Untersee. *Schriften des Vereins für Geschichte des Bodensees und seiner Umgebung* 133: 288-310.

Ostendorp W, Brünner H, Dienst M, Eckmann R, Fiebrich M, Fiedler W, Gabel F, Grabher M, Heck K, Hermann M, Jacoby H, Kiechle J, Knötzsch G, Kramer I, Lehmann A, Löderbusch W, Miler O, Müller M, Rieder J, Sättele B, Schäfer C, Schmidt B, Schmitz G, Schürenberg B, **Stoll S**, Trautner J, Wilhelm P, Winkler U (2014) Ökologische Auswirkungen von Ufermauern und Uferaufschüttungen am Bodensee-Untersee: Ergebnisse einer Expertenbefragung. *Mitteilungen der Thurgauischen Naturforschenden Gesellschaft* 67: 85-105.

+Breyer P, Haase P, Winkelmann C, **Stoll S**, (2013) Skalenabhängiger Einfluss der Gewässerstrukturgüte auf die ökologische Qualität des Makrozoobenthos - Hinweise für die Naturschutzpraxis (Influence de la qualité de la structure des cours d'eau sur la richesse du macrozoobenthos en fonction des échelles spatiales - conseils pour la mise en place de travaux de revitalisation). *Natur + Landschaft INSIDE* 1/13: 29-33.

+Breyer P, Haase P, Winkelmann C, **Stoll S**, (2013) Skalen-abhängiger Einfluss der Gewässerstrukturgüte auf die ökologische Qualität des Makrozoobenthos - Hinweise für die Renaturierungspraxis. *Deutsche Gesellschaft für Limnologie (DGL) – Tagungsbericht 2012 (Koblenz)*: 18-22.

Stoll S, *Früh D, Seeland A, Müller R, Haase P (2013) Der Einfluss des Klimawandels auf aquatische Schnecken: Interaktive Effekte multipler Stressoren Deutsche Gesellschaft für Limnologie (DGL) – Tagungsbericht 2012 (Koblenz): 283-287.

+Früh D, **Stoll S**, Haase P (2012) Sind degradierte Fließgewässerhabitate einem erhöhten Invasionsrisiko durch Neozoen ausgesetzt? Deutsche Gesellschaft für Limnologie (DGL) – Tagungsbericht 2011 (Freising): 192-196.

Sundermann A, Lorenz A, **Stoll S**, Haase P (2010) Renaturierung von Fließgewässern – Hinweise auf dem Weg zum Erfolg. Madgeburger Wasserwirtschaftliche Hefte 10:135-141.

Gabel F, **Stoll S**, Garcia X-F, Pusch M, Fischer P (2009) Schiffs- und windinduzierte Wellen beeinflussen die Nahrungsaufnahme juveniler Cypriniden. Deutsche Gesellschaft für Limnologie (DGL) – Tagungsbericht 2008 (Konstanz): 338-342.

Beeck P, Lages E, **Stoll S**, Hofmann H, Eckmann R, Jatteau P (2009) Der Einfluss von Oberflächenwellen und Bodensubstrat auf die Ei- und Larvenentwicklung von Maifischen *Alosa alosa*. Deutsche Gesellschaft für Limnologie (DGL) – Tagungsbericht 2008 (Konstanz): 329-333.

Stoll S, Fischer P (2006) Die perfekte Welle: Mesokosmosexperimente zur Wirkung von Wellenschlag auf das Wachstum juveniler Cypriniden. Deutsche Gesellschaft für Limnologie (DGL) – Tagungsbericht 2006 (Dresden): 145-149.

Book chapters:

Haase P, Hering D, Hoffmann A, Müller R, Nowak C, Pauls S, **Stoll S**, Straile D (2012) Auswirkungen auf limnische Lebensräume, p91-105. In: Mosbrugger V, Brasseur G, Schaller M, Stribrny B (eds) Klimawandel und Biodiversität - Folgen für Deutschland. Wissenschaftliche Buchgesellschaft, Darmstadt, Germany, 432pp.